

Referenzen

1. Mudge AM, McRae P, McHugh K, et al. Poor mobility in hospitalized adults of all ages. *J Hosp Med*. 2016;11(4):289-291. doi:10.1002/jhm.2536
2. Brown CJ, Redden DT, Flood KL, Allman RM. The Underrecognized Epidemic of Low Mobility During Hospitalization of Older Adults. *J Am Geriatr Soc*. 2009;57(9):1660-1665. doi:10.1111/j.1532-5415.2009.02393.x
3. Abeles, A. *et al.* (2017) 'The relationship between physical activity and post-operative length of hospital stay: A systematic review', *International Journal of Surgery*, 44, pp. 295–302. Available at: <https://doi.org/10.1016/j.ijsu.2017.06.085>.
4. Fazzini, B. *et al.* (2023) 'The rate and assessment of muscle wasting during critical illness: a systematic review and meta-analysis', *Critical Care (London, England)*, 27(1), p. 2. Available at: <https://doi.org/10.1186/s13054-022-04253-0>.
5. Agmon M, Zisberg A, Gil E, Rand D, Gur-Yaish N, Azriel, M. Association Between 900 Steps a Day and Functional Decline in Older Hospitalized Patients. *Jama Intern Med*.2017;177(2):272.274.doi:10.1001/jamainternmed.2016.7266